



Vector spaces and Fourier theory

2 hours

Answer **four** questions. If you answer more than four questions, only your best four will be counted.

- 1 (i) Let  $V$  be a vector space, and let  $v_1, \dots, v_n \in V$ . Define what it means for  $v_1, \dots, v_n$  to
- (a) be linearly independent;
  - (b) span  $V$ ;
  - (c) form a basis for  $V$ . (6 marks)

- (ii) Let  $V = \{f \in C^\infty(\mathbb{R}) : f + f'' = 0\}$  and  $W = \{f \in C^\infty(\mathbb{R}) : f' + f''' = 0\}$ . You may assume that  $V$  and  $W$  are subspaces of  $C^\infty(\mathbb{R})$ .

- (a) Let  $f \in V$ , let  $a = f'(0)$  and let  $b = f(0)$ . Let

$$g(x) = f(x) - a \sin(x) - b \cos(x)$$

and let

$$h(x) = g(x)^2 + g'(x)^2.$$

Show that  $g \in V$  and that  $g(0) = g'(0)$ . Show also that  $h(0) = 0$  and  $h' = 0$ . Deduce that  $h = 0$  and that  $g = 0$ . Hence show that  $\sin$  and  $\cos$  span  $V$ . (11 marks)

- (b) Show that  $V \subseteq W$  and that if  $f \in W$  then  $f' \in V$ . (2 marks)
- (c) Let  $j : \mathbb{R} \rightarrow \mathbb{R}$  be the constant function such that  $j(x) = 1$  for all  $x \in \mathbb{R}$ . Show that  $j \in W$  and that  $\sin$ ,  $\cos$  and  $j$  form a basis for  $W$ . (6 marks)

**2** Let  $V$  be a vector space and let  $U$  and  $W$  be subspaces of  $V$ .

(i) Show that  $U \cap W$  is a subspace of  $V$ . *(5 marks)*

Define the sum  $U + W$  and show that  $U \subseteq U + W$  and  $W \subseteq U + W$ .

*(3 marks)*

(ii) Write down, without proof,

(a) an inequality relating  $\dim U$  and  $\dim V$ ; *(1 mark)*

(b) a strong conclusion about  $U$  and  $V$  whenever  $\dim(U) = \dim(V)$ ; *(1 mark)*

(c) a formula relating  $\dim(U + W)$ ,  $\dim(U \cap W)$ ,  $\dim(U)$  and  $\dim(W)$ . *(1 mark)*

(iii) Suppose that  $\dim(V) = 3$ , that  $\dim U = \dim W = 2$  and that  $U \neq W$ . Show that

(a)  $U + W = V$ ; *(4 marks)*

(b)  $\dim(U \cap W) = 1$ . *(2 marks)*

(iv) Let

$$V = \{(a, b, c, d)^T \in \mathbb{R}^4 \mid a + b + c + d = 0\}$$

$$U = \{(a, b, c, d)^T \in \mathbb{R}^4 \mid a + b + c + d = 0 = a + b\} \text{ and}$$

$$W = \{(a, b, c, d)^T \in \mathbb{R}^4 \mid a + b + c + d = 0 = a - c\}.$$

You may assume that these are all subspaces of  $\mathbb{R}^4$ .

(a) Write down, without proof,  $\dim(V)$ ,  $\dim(U)$  and  $\dim(W)$ . Find a vector  $v \in \mathbb{R}^4$  that spans  $U \cap W$ . *(4 marks)*

(b) Let  $u = (2, -2, 7, -7)$  and let  $w = (3, -5, 3, -1)$ . Find  $u' \in U$  and  $w' \in W$  such that  $u' + w' = u + w$  but  $u' \neq u$ . *(4 marks)*

- 3 (i) Let  $V$  and  $W$  be vector spaces and let  $\phi : V \rightarrow W$  be a linear map. Define the *kernel*  $\ker(\phi)$  and the *image*  $\text{im}(\phi)$ . **(2 marks)**

State, without proof, a formula relating  $\dim(\ker(\phi))$ ,  $\dim(\text{im}(\phi))$  and  $\dim(V)$ . **(1 mark)**

- (ii) Let  $V = M_2(\mathbb{R})$  be the vector space of  $2 \times 2$  real matrices, let

$$A = \begin{pmatrix} 0 & 1 \\ 2 & 3 \end{pmatrix}$$

and let  $\phi : V \rightarrow V$  be the map such that  $\phi(X) = AX - XA$  for all  $X \in V$ . You may assume that  $\phi$  is linear.

- (a) Find the matrix of  $\phi$  with respect to the basis

$$E_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, E_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, E_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, E_4 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

**(8 marks)**

- (b) Show that  $I_2 \in \ker(\phi)$  and that  $A \in \ker(\phi)$ . Deduce that  $\dim(\ker(\phi)) \geq 2$  and  $\dim(\text{im}(\phi)) \leq 2$ . **(5 marks)**

Write down two linearly independent elements of  $\text{im}(\phi)$  and show that  $\dim(\text{im}(\phi)) = 2 = \dim(\ker(\phi))$ . **(3 marks)**

- (c) Let  $W$  be the vector space of  $2 \times 2$  real matrices with trace 0. Show that  $\text{im}(\phi) \subseteq W$ . Is  $\text{im}(\phi) = W$ ? Justify your answer. **(3 marks)**

- (d) Show that  $A^2 \in \ker(\phi)$  and express  $A^2$  as a linear combination of  $I_2$  and  $A$ . **(3 marks)**

- 4 (i) Define the notion of an *inner product* on a finite-dimensional vector space over  $\mathbb{R}$ . **(5 marks)**

- (ii) Let  $V$  be an inner product space (over  $\mathbb{R}$ ). State the Cauchy-Schwarz inequality for vectors  $v, w \in V$ , including the criterion for when equality holds. **(3 marks)**

- (iii) Consider the inner product space  $C[0, 1]$  with inner product given by

$$\langle f, g \rangle = \int_0^1 f(x)g(x) dx.$$

- (a) Show that for any  $f \in C[0, 1]$ , we have

$$\left| \int_0^1 (1 - x^3)f(x) dx \right| \leq \frac{3}{\sqrt{14}} \sqrt{\int_0^1 f(x)^2 dx}. \quad \textbf{(6 marks)}$$

- (b) Use the Gram-Schmidt process to find an orthogonal basis for the subspace of the inner product space  $C[0, 1]$  spanned by  $1, x$  and  $x^4$ . **(11 marks)**

- 5 (i) Let  $V$  be a vector space with an inner product and let  $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$  be a finite subset of  $V$ . Explain what it means for  $\mathcal{V}$  to be *orthogonal* and what it means for  $\mathcal{V}$  to be *strictly orthogonal*. **(3 marks)**

Show that if  $\mathcal{V}$  is strictly orthogonal then  $v_1, v_2, \dots, v_n$  are linearly independent. **(5 marks)**

- (ii) Consider the Fourier inner product space  $C[-\pi, \pi]$  of continuous functions  $[-\pi, \pi] \rightarrow \mathbb{R}$  with the inner product

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f(t)g(t) dt.$$

Let  $f(t) = \sin t \cos 5t$  and  $g(t) = \sin 2t \cos 4t$ . Calculate the angle between  $f$  and  $g$  and the distance between  $f$  and  $g$ . **(11 marks)**

- (iii) Consider the subspace  $V$  of the Fourier inner product space of trigonometric polynomials of degree at most 1, spanned by the set  $\mathcal{V} = \{1, \cos t, \sin t\}$ , and the space  $M_2(\mathbb{R})$  of  $2 \times 2$  real matrices with inner product  $\langle A, B \rangle = \text{trace}(A^T B)$ . If  $v = \alpha + \beta \cos t + \gamma \sin t$ , define  $\phi(v) = \begin{pmatrix} \beta & \alpha \\ 0 & \gamma \end{pmatrix}$ .

If  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , compute  $\langle \phi(v), A \rangle$ , and hence give the adjoint map  $\hat{\phi} : M_2(\mathbb{R}) \rightarrow V$ . **(6 marks)**

**End of Question Paper**